

Article

Cite this article: Takasaki, R., Y. Kobayashi, A. R. Fiorillo, T. Chinzorig, and G. F. Funston (2026). Gastrolith shape as an indicator of digestive function and its implications for dinosaurian digestive strategies. *Paleobiology*, 1–12. <https://doi.org/10.1017/pab.2026.10112>

Received: 17 July 2025 UTC

Revised: 02 June 2026 UTC

Accepted: 09 June 2026 UTC

Handling Editor:

Alexander Dunhill

Corresponding author:

Ryuji Takasaki;

Email: r-takasaki@ous.ac.jp

Gastrolith shape as an indicator of digestive function and its implications for dinosaurian digestive strategies

Ryuji Takasaki¹ , Yoshitsugu Kobayashi², Anthony R. Fiorillo³, Tsogtbaatar Chinzorig^{4,5} and Gregory F. Funston⁶

¹Faculty of Biosphere-Geosphere Science, Okayama University of Science, Okayama, Japan

²Hokkaido University Museum, Hokkaido, Japan

³New Mexico Museum of Natural History & Science, Albuquerque, New Mexico, U.S.A.

⁴Department of Biological Sciences, North Carolina State University at Raleigh, U.S.A.

⁵Division of Paleozoology, The Institute of Paleontology, Mongolian Academy of Sciences, Mongolia

⁶Department of Anatomical Sciences, Renaissance School of Medicine, Stony Brook University, U.S.A.

Abstract

The evolution of the muscular stomach was a key innovation in the avian body plan, enabling high metabolic rates by functionally replacing oral processing. However, its deep-time origins within Archosauria have remained poorly understood, primarily due to the extreme rarity of fossilized stomachs. Here, we demonstrate that gastrolith shape is a robust proxy for digestive function and, by extension, stomach muscularity and diet. By assembling a comprehensive dataset from 104 individuals across 46 extant archosaur species, we establish a strong correlation between gastrolith shape, stomach muscularity, and diet. Application of this validated method to the fossil record reveals divergent digestive strategies among major dinosaur clades. Herbivorous ornithischians retained angular gastroliths, consistent with limited gastric abrasion and compensation through oral processing. Sauropods likewise retained angular gastroliths, consistent with limited gastric abrasion and digestion relying more on long retention times and fermentation. Conversely, the muscular stomach, indicated by rounded gastroliths, evolved early within Theropoda, appearing at least by the base of Maniraptoriformes. This innovation was likely a crucial prerequisite for repeated transitions to herbivory in maniraptoriform theropods with reduced dentition, shifting mechanical processing from the jaws to the stomach and reducing reliance on heavy jaw adductor musculature.

Non-technical Summary

For decades, paleontologists have often interpreted the discovery of stomach stones, or gastroliths, inside a dinosaur's ribcage as a sign of a plant-based diet. However, because many modern carnivores also swallow stones, this assumption has been uncertain. This study reveals that the secret to understanding a dinosaur's digestion lies not in the presence of stones, but in their shape. Analysis of a new, comprehensive dataset from modern birds and crocodiles shows that rounded stones are a clear indicator of a muscular stomach used to grind tough, fibrous plants. Angular stones, in contrast, point to a weaker stomach and reduced gastric processing. Applying this new lens to the fossil record reveals a major evolutionary divergence. Iconic herbivores like the distant relatives of *Triceratops* retained angular stones, consistent with limited gastric abrasion and reliance on chewing and other processing. Long-necked sauropods also retained angular stones, suggesting limited gastric abrasion. Their digestion likely relied more on retention time and fermentation rather than a powerful grinding stomach. In contrast, the lineage of meat-eating dinosaurs that led to birds evolved muscular gizzards early in their history. This key innovation allowed them to process tough foods without relying on heavy jaw musculature, paving the way for subsequent skull and feeding adaptations in the avian stem lineage.

Introduction

Birds mechanically process food in a muscular stomach (the gizzard), sometimes aided by retained gastroliths. In domestic geese (*Anser anser*), the gizzard serves as the primary site of mechanical food breakdown (Moore 1998). This process is sometimes aided by ingested stones (gastroliths) retained in the gizzard (Wings 2007). This combination of muscles and stones can achieve food processing efficiencies comparable to those of the highly complex teeth of herbivorous mammals (Fritz et al. 2011). Because gastroliths are especially common among herbivorous birds (Gionfriddo and Best 1999), their presence has been considered evidence of herbivory in some dinosaurs (Kobayashi et al. 1999; Zanno and Makovicky 2011), particularly if the animal

© The Author(s), 2026. Published by Cambridge University Press on behalf of Paleontological Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

PALEOBIOLOGY
A PUBLICATION OF THE
 **Paleontological SOCIETY**

 **CAMBRIDGE**
UNIVERSITY PRESS



hosts a large number of gastroliths, often found as a tight cluster within the abdominal region, that comprise more than 1% of body mass (Wings and Sander 2007). This gastrolith–herbivory relationship is now widely used by vertebrate paleontologists to infer herbivory in several dinosaur lineages (Cerde 2008; Makovicky et al. 2011; Zanno and Makovicky 2011).

However, large numbers of gastroliths are also common in insectivorous (Barlow et al. 1963; Jenkinson and Mengel 1970; Brown 1976; Barrentine 1980; Potter 1983), piscivorous (Lambrecht 1933; McKeown 1934; Siegel-Causey 1990; Beaune et al. 2009), and even carnivorous birds (Röhrig 1907; Lambrecht 1933), raising doubts that the gastrolith–herbivory relationship is as straightforward as presumed. Indeed, Takasaki and Kobayashi (2020b) demonstrated that the relationship between body mass and gastrolith mass in strictly carnivorous Crocodylia, the sister group of Dinosauria, is statistically indistinguishable from that of herbivorous birds. Additionally, a hypercarnivorous dinosaur, *Tarbosaurus* (MPC-D 552-1), also possessed abundant gastroliths (Fig. 1C). This ambiguity is also evident in Early Cretaceous birds: abdominal stone masses initially discussed as gastroliths in relation to herbivory or diet switching in *Yanornis* (Zhou et al. 2004) were later interpreted as likely representing sand impacted in the intestines (Zheng et al. 2014). Similarly, structures initially described as a gastrolith mass in *Bahaiornis* (Li et al. 2014) were subsequently interpreted as postmortem mineral precipitation (Liu et al. 2021). In light of such taphonomic and interpretive uncertainties, the presence of gastrolith-like stone accumulations should be treated cautiously when inferring diet in stem birds (O'Connor 2019) and recent multiproxy frameworks for fossil-bird diet reconstruction emphasize synthesizing independent evidence rather than relying on any single indicator (Miller and Pittman 2021).

These issues motivate evaluation of an alternative approach: rather than treating gastrolith occurrence as diagnostic, gastrolith shape may record the mechanical environment of the stomach and

thereby provide an independent line of evidence for multiproxy dietary inference. Consistent with this possibility, Takasaki and Kobayashi (2020a) experimentally demonstrated in domestic chicks (*Gallus gallus domesticus*) that dietary differences can generate significant differences in gastrolith shape: whereas plant feeders have rounded gastroliths, those of vertebrate feeders are rougher and more angular. This pattern is driven by gizzard muscularity, a plastic trait that develops in response to diet, because gastrolith shape differences stem from intragastric abrasions (Wings and Sander 2007; Takasaki and Kobayashi 2020a), which in turn rely on mechanical action. Indeed, chicks with higher gizzard muscularity generally have more rounded gastroliths than those with lower gizzard muscularity (Takasaki and Kobayashi 2020a). Although Wings (2007) warned against using gastrolith morphology to predict function, the recent experiment (Takasaki and Kobayashi 2020a) demonstrated that gastrolith morphology reflects the degree of gastric mechanical digestion, at least in domestic chicks under experimental conditions, opening a window toward further investigations into whether the same trend can be observed among wild archosaurs, and furthermore, whether the method can be applied to extinct taxa.

Here we build a comprehensive dataset of gastrolith shape, stomach muscularity, and diet from a wide range of extant archosaurs (crocodylians and birds) to investigate whether the trends observed in domestic chicks—that gastrolith shape reflects diet and stomach muscularity—are broadly applicable across diverse, non-experimental archosaur taxa. Using the empirical patterns validated in extant species, we analyze the shapes of the gastroliths of the extinct taxa to infer the relationships between gastrolith shape, diet, and stomach muscularity throughout the evolutionary history of Archosauria. This project sheds light on the evolutionary history of the archosaur gastrointestinal system, which is currently largely unknown due to the extremely limited fossil record (Dal Sasso and Maganuco 2011; Wang et al. 2022).




Figure 1. Close-up views of dinosaur gastroliths (A–D). A, *Haya griva* (IGM 100/2015); B, *Archaeorhynchus spatula* (IVPP V17091); C, *Tarbosaurus bataar* (MPC-D 552-1); D, close-up of *Tarbosaurus* gastrolith mass showing the red boxed region of C. Red arrowheads are pointing to the major (not all) gastroliths in the abdominal cavity. E, The five categories used to characterize gastrolith shape, redrawn from Best and Gionfriddo (1991). See Supplementary Figs. S1–S3 for the other specimens analyzed in this study.

Materials and Methods

Specimens

To test the relationships between gastrolith shape, diet, and stomach muscularity, we examined 382 extant specimens, from 104 of which we could collect at least one gastrolith, representing 42 neornithine (90 individuals) and 4 crocodylian (14 individuals) taxa (Supplementary Data S1, S2, Supplementary Table S6). Of these, 56 individuals meeting the ≥ 35 gastrolith threshold (see “Results”) were used for model training and subsequent analyses. The majority of the neornithine specimens are accessioned at the Hokkaido University Museum (HoUMVC) or the Botanic Garden of Hokkaido University (HUNHM) collections and are publicly accessible. Two birds without official collection numbers (Bird 0001 and Bird 0002) are also deposited at the Hokkaido University Museum and are publicly available, but they are not associated with any skeletal materials. Four crocodylian specimens (Darwin 01–04) are housed at the Crocodylus Park, NT, Australia. The others (croc 01–10) were provided by Koike Wani Sohono Company (Shizuoka Prefecture, Japan) and Atagawa Tropical & Alligator Garden (Shizuoka Prefecture, Japan), and the gastroliths are deposited at the Hokkaido University Museum, but no skeletal elements are associated to them. All neornithines were assigned to dietary categories proposed in Wilman et al. (2014), which are assigned based on their summed scores of constituent individual diets: VertFishScav, PlantSeed, Invertebrate, and Omnivore. These broad, standardized categories provide consistent training labels across a wide extant taxonomic sample and are appropriate for a supervised model designed to be applied across Archosauria. All crocodylians were considered VertFishScav. Stomach muscularity was calculated as the ratio of stomach mass to body mass (Takasaki and Kobayashi 2020a).

Gastroliths of 29 extinct archosaur specimens were examined (Supplementary Data S3). All of the gastroliths subject to the analyses are strictly limited to the stones found in the abdominal regions of the host animal. Therefore, the present analyses have no a priori assumption based on morphology in identifying gastroliths, following the suggestion by Wings (2007). Note that this does not necessarily indicate that the stones were preserved within the stomach or “gizzard” in life.

Gastrolith Shape Evaluations

Gastrolith shapes in extant neornithines and crocodylians were evaluated quantitatively and qualitatively. The quantitative method follows the experiment performed by Takasaki and Kobayashi (2020a): Circularity, Roundness, and Solidity of the extant archosaur gastroliths larger than 0.5 mm in minor axis are evaluated using ImageJ v. 1.8.0 (Schneider et al. 2012) (Supplementary Data S1). While the quantitative method relies completely on two-dimensional information, an experiment on domestic chickens demonstrated that the quantitative proxy successfully differentiates plant feeders from vertebrate feeders (Takasaki and Kobayashi 2020a). As for the qualitative evaluation, gastroliths were categorized following Best and Gionfriddo (1991) into five categories: Angular, Sub-angular, Sub-rounded, Rounded, and Well-rounded (Fig. 1E, Supplementary Data S2). Following the original description by Best and Gionfriddo (1991), the five categories are defined as follows: Angular, sharp and irregular corners; Sub-angular, corners slightly rounded and inlets sharp; Sub-rounded, corners rounded and inlets more or less smooth; Rounded, corners well-rounded and only a few inlets; and Well-rounded, smoothly rounded with no corners or inlets. While the qualitative proxy could

potentially be subjective, qualitative stone shape evaluation has proven to be useful in field geology for nearly a century (Russell and Taylor 1937; Powers 1953). Furthermore, the qualitative categories integrate corner sharpness and concavities (inlets) defined by Best and Gionfriddo (1991), which were evaluated from multiple angles. In order to be consistent, R.T. evaluated the gastrolith shapes. The two proxies are thus expected to complement the weaknesses of each other.

Multiple mechanisms can contribute to a nonrandom distribution of pre-consumption gastrolith shapes, including shape-based selectivity by the host animal (Best and Gionfriddo 1991, 1994; Takasaki and Kobayashi 2020a) and the depositional environment of the source sediment (e.g., riverbanks, pebbly beaches, or weathered conglomerates). The relative contribution of these factors to the shape distribution at the moment of ingestion is intrinsically unknown for both the extant and fossil specimens examined here—for the extant birds, rearing or capture locality (Supplementary Table S6) does not necessarily indicate where the gastroliths were acquired. This study treats this as an unavoidable limitation of any gastrolith-based study and tests whether, despite such pre-consumption variation, gastrolith shape nonetheless tracks diet and stomach muscularity in our extant training set; the empirical adequacy of treating gastrolith shape as a proxy for these variables is evaluated in the “Discussion.”

Because dinosaur gastroliths often remained embedded within the matrix for conservation or exhibition purposes, the quantitative proxy could not be used, as the method requires gastroliths to be completely isolated (Takasaki and Kobayashi 2020a). Therefore, dinosaur gastrolith shapes were evaluated using only the qualitative proxy. Because not all gastroliths are exposed on the surface, only the exposed ones were evaluated. This sampling strategy is intended to represent the original gastrolith assemblage, but it relies on the working assumption that postmortem processes did not cause systematic shape-dependent sorting within the carcass. Generally, all gastroliths that are exposed and are larger than 0.5 mm in minor axis were evaluated, but in some cases, specimens contained far more than 1000 gastroliths, which made it impractical to evaluate every gastrolith. In those rare cases, approximately 1000 randomly selected gastroliths were evaluated. A sensitivity test indicated that the 95% interval of subsample means converges well below 1000 stones ($n \approx 100$ –150 for ± 0.2 SD; $n \approx 500$ –520 for ± 0.1 SD, where SD is the within-specimen variation), suggesting that scoring ~ 1000 stones is sufficient to represent the exposed population. Among the 29 specimens subject to the analyses, gastrolith shapes were evaluated based on direct observations (23 specimens) or high-resolution photographs (6 specimens) focused on the gastroliths (Supplementary Data S3). Published figures from the literature were, in most cases, insufficient for capturing subtleties of gastrolith shape that are critical for categorizing stone morphotype. Where published figures were insufficient, additional images were requested when feasible; specimens were excluded when adequate resolution could not be obtained, which likely reduces taxonomic coverage but minimizes misclassification risk.

Statistical Analyses

To test whether the quantitative gastrolith shape reflects avian diet as in domestic chicks (Takasaki and Kobayashi 2020a), discriminant analyses were conducted for both quantitative and qualitative proxies. Before the analyses, quantitative shape proxies (i.e., Circularity, Roundness, Solidity) were averaged per individual and the ratios of the qualitative gastrolith shape categories (i.e., Angular, Sub-angular, Sub-rounded, Rounded, Well-rounded) were arcsine-transformed. This transformation is commonly applied to proportion data bounded

between 0 and 1 to reduce heteroscedasticity and improve approximate normality for methods (Zar 2010). To test the sensitivity of the discriminant accuracy against gastrolith number, 1–50 gastroliths were subsampled from each individual and discriminant analyses were repeated 100 times. The minimal number of gastroliths to reach equilibrium in discriminant accuracy was inferred from this sensitivity test. Subsequent analyses were performed on a pruned dataset of specimens with more gastroliths than the point of equilibrium.

Principal component analyses (PCA) were performed on the pruned datasets to reduce the dimensionality of the quantitative and qualitative shape indices. Standard major axis (SMA) regression was then performed to test the relationship between gastrolith shape and stomach muscularity. Correlation between the first principal axes of the quantitative and the qualitative proxies was also tested using SMA to verify that the qualitative proxy adequately reflects the quantitative proxy. Following this validation, the diet prediction model was trained by performing linear discriminant analysis on the pruned qualitative dataset. Extinct archosaur diets were predicted by applying the prediction model. Note that predicted diets are necessarily expressed using the EltonTraits categories (Wilman et al. 2014), because the classifier model is trained on those labels. The evolution of gastrolith shape, which reflects stomach muscularity (Takasaki and Kobayashi 2020a), was inferred from ancestral-state reconstruction. The ancestral-state reconstruction was performed based on a composite phylogenetic framework of Boyd (2015), Sander et al. (2011), Hendrickx et al. (2015), and O'Connor (2019). The tree is used as a scaffold for trait mapping and exploratory ancestral-state reconstruction and is not intended as a new phylogenetic estimate with newly inferred support values. The composite topology was assembled to reflect widely used higher-level relationships among major archosaur clades based on the cited sources, and contentious nodes are not interpreted beyond their influence on broad-scale patterns. Because phylogenetic uncertainty is not explicitly modeled here, ancestral-state reconstructions are treated as exploratory and conditional on the adopted topology. The framework is provided as [Supplementary Data S4](#). All analyses were performed on R v. 4.4.1 (R Core Team 2020), using R packages *ape* v. 5.8 (Paradis and Schliep 2019), *MASS* v. 7.3-60.2 (Venables and Ripley 2013), *geiger* v. 2.0.11 (Pennell et al. 2014), and *smatr* v. 3.4-8 (Warton et al. 2012).

Terminology

The stones considered in this study are, strictly speaking, “geogastroliths” (swallowed sediment particles of no caloric value retained in the digestive tract of an animal), following the classification by Wings (2007). However, for brevity and readability, the simpler term “gastrolith” is used consistently throughout this paper to refer to these objects. This general term is chosen over more specific ones like “stomach stone” or “gizzard stone” for two main reasons. First, while the gastroliths from extant specimens were collected directly from the stomach, the precise anatomical location of stone masses in fossils cannot be definitively determined. “Gastrolith” is therefore a more accurate term for fossil specimens and is used for all samples to maintain consistency. Second, and more importantly, terms like “gizzard stone” carry a strong functional presumption of mechanical grinding, although such function is often absent in carnivorous birds (Wings 2007). Furthermore, the term “gizzard stone,” common in ornithology, presupposes the presence of an avian-like gizzard. However, because the fundic stomach of extant crocodylians (often regarded as a “gizzard”) is not homologous with the avian gizzard (Takasaki and Kobayashi

2020b), and thus the timing of true gizzard acquisition is unknown, using the term “gizzard stone” would be presumptive within an evolutionary context. The neutral term “gastrolith” avoids such locational, functional, and developmental presumptions.

For similar reasons, the index for stomach muscle development is termed “stomach muscularity” (defined as the ratio of ventriculus mass to body mass) rather than “gizzard muscularity,” which is the term used by Takasaki and Kobayashi (2020a). This avoids assuming the presence of a true avian-like gizzard (muscular ventriculus), an organ homologous to the pyloric stomach of crocodylians (Takasaki and Kobayashi 2020b). Crocodylian stomach muscularity was calculated by the ratio of the fundic stomach mass to the body mass.

For readability, the following text labels are used for the Elton-Traits diet categories (Wilman et al. 2014): “VertFishScav” = Vertebrate Feeders (including fish and scavenging), “PlantSeed” = Plant and Seed Feeders, “Invertebrate” = Invertebrate Feeders, and “Omnivore” = Omnivores. Capitalized category labels (e.g., Vertebrate Feeders) refer specifically to these EltonTraits training labels and model outputs. For general comparisons and descriptions of extinct archosaur diets, lowercase dietary terms are used (e.g., herbivorous, granivorous, invertivorous, vertivorous) following the glossary compiled by Miller and Pittman (2021), preferentially using inclusive terms unless narrower specializations are independently supported. Accordingly, this study interprets the Elton-Traits categories as coarse equivalents for comparison with prior inferences: Plant and Seed Feeders to herbivory, Vertebrate Feeders to vertivory, Invertebrate Feeders to invertivory, and Omnivores to omnivory.

Institutional Abbreviations. AMNH: American Museum of Natural History, New York, NY, USA; DMNH, Denver Museum of Nature & Science, Denver, Colorado, USA; GSGM: Gansu Geological Museum, Lanzhou, Gansu, China; HUNHM: Hokkaido University Natural History Museum (Botanic Garden & Museum, Hokkaido University), Sapporo, Hokkaido, Japan; HoUM: Hokkaido University Museum, Sapporo, Hokkaido, Japan; IGM: Mongolian Institute of Geology, Ulaanbaatar, Mongolia; IVPP: Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, Beijing, China; MPC-D: Mongolian Paleontological Center, Dinosaur Collection, Ulaanbaatar, Mongolia; NMMNH: New Mexico Museum of Natural History and Science, Albuquerque, New Mexico, USA; STM: Shandong Tianyu Museum of Nature, Pingyi, Shandong, China.

Results

The sensitivity tests on minimal stomach stone counts indicate that the discriminant accuracies steadily improve with an increasing number of subsampled stomach stones for both qualitative and quantitative proxies (Fig. 2A,B). The discriminant accuracies reached an equilibrium at approximately 35 stomach stones in both proxies, at which point the overall accuracy is approximately 75% in the qualitative proxy and approximately 65% in the quantitative proxy. Considering this result, only individuals with at least 35 stomach stones were used for the subsequent analyses to maximize the discriminant accuracy. Applying the ≥ 35 stone threshold pruned 56% of Plant and Seed Feeders, 69% of Vertebrate Feeders, 17% of Invertebrate Feeders, and 39% of Omnivores, leaving 56 individuals for subsequent analyses (15 Plant and Seed Feeders, 17 Vertebrate Feeders, 20 Invertebrate Feeders, and 4 Omnivores).

The final discriminant analysis using the pruned training dataset resulted in an overall accuracy of 78.57% for qualitative proxy and




Figure 2. Relationships among gastrolith shape, diet, and stomach musculature. Transition of discriminant accuracy along increase in subsampling size in qualitative (**A**) and quantitative (**B**) proxies. Note that the overall accuracy (black) reaches the equilibrium at around $n = 35$. Bivariate plots of linear discriminant 1 (LD 1) and LD 2 scores obtained from the linear discriminant analyses in qualitative (**C**) and quantitative (**D**) proxies. Inner and outer ellipses represent 50% and 95% confidence for each diet category, respectively. Standard major axis (SMA) regression plot of gastrolith shape (principal component 1 [PC 1]) on stomach musculature of extant archosaurs in qualitative (**E**) and quantitative (**F**) proxies. The gray region represents the 95% confidence interval.

67.86% for the quantitative proxy (Supplementary Table S1). The discriminant accuracies varied considerably among dietary categories: Plant and Seed Feeders had the highest accuracy (qualitative: 86.67%, quantitative: 86.67%), followed by Invertebrate Feeders (qualitative: 85.00%, quantitative: 75.00%) and Vertebrate Feeders (qualitative: 76.47%, quantitative: 58.82%), and Omnivores had the lowest accuracy (qualitative: 25.00%, quantitative: 0.00%).

For the qualitative proxy, intergroup variance was best represented by the proportions of Angular and Well-rounded stomach stones (linear discriminant axis 1 [LD 1]: 75.51%; Supplementary Table S3), followed by Sub-angular and Rounded stomach stones (LD 2: 18.62%). Similarly, principal component analysis shows that the largest source of shape variation was the proportions of Angular, Rounded, and Well-rounded stomach stones (principal component 1 [PC 1]: 74.32%), followed by proportions of Sub-angular and Sub-rounded stomach stones (PC 2: 14.57%). For the quantitative proxy, LD 1 (88.04%) was primarily represented by Circularity and Solidity, while Roundness had a high loading on LD 2 (10.23%). The corresponding principal component analysis shows that PC 1 (53.02%) was driven by Circularity and Solidity, while PC 2 (32.73%) was driven by Roundness.

PC 1 values derived from the qualitative shape proxies correlated well with those from the quantitative proxy ($p < 0.001$, $R^2 = 0.66$). For both proxies, the PC 1 scores also correlated significantly with the stomach muscularity ($p < 0.001$, $R^2 = 0.44$ for qualitative and $R^2 = 0.37$ for quantitative proxy).

Building on the extant training set, Elton Trait diet categories of 29 dinosaur specimens (16 taxa) were predicted based on preserved stomach stones (Fig. 2C, Table 1; see Supplementary Text S1 for detailed descriptions of dinosaur stomach stones). Because dinosaur gastroliths are often embedded in the matrix, only the qualitative model was utilized for this analysis (see “Materials and Methods” and “Discussion” for validations). Note that although extant bird specimens with fewer than 35 stomach stones were pruned in the training dataset, fossil specimens with low stone counts were still analyzed. The specimens with low stomach stone count (*Iteravis*, IVPP V18958; and *Jeholornis*, STM 3-19; stomach stone $n = 6$ each) were recovered far from the morphospace occupied by all other specimens (Supplementary Table S5). The predicted dietary habits of dinosaurs with more than 35 stomach stones are generally concordant with the previous results (Table 1), with the notable exceptions of several ornithischians (*Haya*, *Psittacosaurus*), a sauropod (*Cedarosaurus*), and several Mesozoic birds (*Bellulornis*, *Gansus*, and *Iteravis*).

The ancestral-state reconstruction of PC 1 stomach stone shape values yielded the following results for key nodes: ancestral Dinosauria was reconstructed with a value of -1.52 , ancestral Saurischia with -1.65 , ancestral Theropoda with -1.63 , ancestral Maniraptoriformes with -2.48 , and ancestral Avialae with -2.62 (Fig. 3).

Discussion

The sensitivity analysis on the minimum stomach stone counts for predicting archosaur diet demonstrates the critical importance of sample size. The discriminant accuracy of our models improved substantially as the number of subsampled stones increased, stabilizing only after approximately 35 stones were included in the analysis (Fig. 2A,B). This finding highlights the necessity of averaging across a large population of stones to mitigate the inherent variability of individual stones. The shape of any single stomach stone is influenced by numerous factors, including its time since ingestion, its original shape, abrasion rate, and its lithological

properties such as hardness (Wings 2007). Analyzing a large population of stones from a single individual permits the effects of these confounding variables to be averaged out, revealing a more robust signal of the overall abrasive environment within the stomach.

However, setting this threshold involves a trade-off. While a higher minimum stone count improves the reliability of each data point, it also reduces the number of specimens available for the training dataset, which can, in turn, decrease the overall predictive power of the model. In the present case, applying the $n = 35$ threshold reduced our extant dataset from 104 to 56 individuals. Therefore, the optimal threshold must be a point that maximizes per-specimen reliability without excessively compromising the training dataset size. This study chose a minimum of 35 stones because this represents the point of equilibrium where discriminant accuracy plateaued for both our qualitative and quantitative proxies, while still retaining a sufficiently large dataset for robust model creation. The data pruning resulted in a relatively even distribution of sample sizes across diet categories, with the exception of omnivores (see “Results”).

Notably, the predictive accuracy of the model based on the qualitative proxy (78.57%) was higher than that of the quantitative proxy (67.86%). Although the two proxies capture similar features, as supported by the correlation between their respective PC 1 values ($R^2 = 0.66$), this difference in accuracy implies they are not identical in the information they retain. This discrepancy may highlight the importance of three-dimensional information. While the quantitative proxy relies solely on two-dimensional outlines from photographs, the qualitative proxy is based on human observation, which can incorporate three-dimensional shape information. The human eye is better at capturing subtle features, such as the small inlets characteristic of the Angular and Sub-angular categories, which may not always be resolved in the two-dimensional photographs used for quantitative analysis. Therefore, our results suggest that the qualitative proxy not only serves as a valid substitute for the quantitative method (Takasaki and Kobayashi 2020a) when dealing with embedded fossils, but may even outperform it in terms of predictive accuracy.

The results of our discriminant and principal component analyses are highly concordant with previous empirical experiments and inferred functions of wild bird stomach stones (Wings 2007). Greater LD 1 values (i.e., a dominance of Angular stomach stones) in Invertebrate Feeders and Vertebrate Feeders compared with Plant and Seed Feeders align with experiments on domestic chicks, which demonstrated rougher, more angular stomach stones in carnivorous groups than in herbivorous groups (Takasaki and Kobayashi 2020a). Likewise, the dominance of rounded stomach stones in specimens with high stomach muscularity, demonstrated by the correlation between PC 1 and stomach muscularity (Fig. 2E,F), is consistent with these same experimentally derived results (Takasaki and Kobayashi 2020a).

This framework reveals that the shape-based analysis is not merely a diet proxy, but more fundamentally, a proxy for mechanical digestive function. The causal chain can be understood as follows: the diet of an animal influences the required stomach muscularity, which in turn dictates the mechanical function of the stomach (e.g., an active grinding mill). This function is what directly influences stomach stone shape through abrasion. Therefore, by observing stone shape, the most direct inference that can be made is about the mechanical function of the stomach. For instance, the prediction of angular stones in some carnivores including *Buteo buteo*, despite having a large number of gastroliths (Supplementary Data S2), suggests the absence of a grinding function, which is consistent with the use of stones for nonabrasive purposes like range or simply an accidental intake (Wings 2007).

Table 1. Predicted diets of extinct archosaurs based on gastrolith shapes. The italic predicted diet represents the predictions that differ from previous assumptions. Note that prediction accuracy is expected to be low on the specimens with fewer gastroliths, even if the posterior probabilities are high

Taxa	ID	Clade	Predicted diet	Previous suggestion	No. of gastroliths	Posterior probability (%)			
						Invertebrate	Omnivore	Plant + Seed	Vertebrate
<i>Archaeorhynchus spathula</i>	IVPP V14287	Avialae	Plant + Seed	Herbivory	38	0	0.3	99.69	0.01
<i>Archaeorhynchus spathula</i>	IVPP V17075	Avialae	<i>Omnivore</i>	Herbivory	64	0.06	74.34	21.16	4.44
<i>Archaeorhynchus spathula</i>	IVPP V17091	Avialae	Plant + Seed	Herbivory	79	0.84	6.21	71.52	21.42
<i>Archaeorhynchus spathula</i>	IVPP V20312	Avialae	<i>Vertebrate</i>	Herbivory	44	1.33	23.15	36.59	38.93
<i>Bellulornis rectusunguis</i>	IVPP V17970	Avialae	<i>Vertebrate</i>	Herbivory	17	0.31	1.25	6	92.43
<i>Caudipteryx dongi</i>	IVPP V12344	Theropoda	Plant + Seed	Herbivory	196	0.02	8.28	91.3	0.4
<i>Caudipteryx sp.</i>	IVPP V12430	Theropoda	Plant + Seed	Herbivory	202	0.01	1.9	98.04	0.05
<i>Cedarosaurus weiskopfae</i>	DMNH 39045	Sauropoda	<i>Vertebrate</i>	Herbivory	96	3.64	2.62	5.7	88.04
<i>Deinococheirus mirificus</i>	MPC-D 100/127	Theropoda	<i>Plant + Seed</i>	Omnivory	959	0	0.16	99.83	0
<i>Diplodocus hallorum</i>	NMMNH 3690	Sauropoda	Plant + Seed	Herbivory	117	0.05	0.08	99.71	0.16
<i>Eogranivora edentulata</i>	STM35-3	Avialae	Plant + Seed	Herbivory	32	0.09	0.04	99.25	0.62
<i>Gansus yumenensis</i>	GSGM 06-CM-011	Avialae	<i>Vertebrate</i>	Herbivory?	34	4.01	15.3	0.34	80.35
<i>Gansus yumenensis</i>	GSGM 07-CM-011	Avialae	<i>Omnivore</i>	Herbivory?	18	0.04	98.14	1.55	0.27
<i>Haya griva</i>	IGM 100/2015	Ornithischia	<i>Omnivore</i>	Herbivory	42	0.88	85.77	2.4	10.95
<i>Haya griva</i>	IGM 100/3182	Ornithischia	<i>Vertebrate</i>	Herbivory	43	2.56	1.6	20.88	74.97
<i>Iteravis huchzermeyeri</i>	IVPP V18958	Avialae	<i>Invertebrate</i>	Herbivory?	6	98.1	0.25	1.65	0
<i>Iteravis huchzermeyeri</i>	IVPP V20133	Avialae	<i>Vertebrate</i>	Herbivory?	17	0.12	3.45	13.4	83.03
<i>Iteravis huchzermeyeri</i>	IVPP V20134	Avialae	<i>Vertebrate</i>	Herbivory?	19	0.13	10.33	24.48	65.06
<i>Jeholornis prima</i>	STM2-15	Avialae	Plant + Seed	Herbivory	49	0.15	3.75	94.93	1.17
<i>Jeholornis prima</i>	STM2-31	Avialae	Plant + Seed	Herbivory	63	0.01	3.54	96.05	0.4
<i>Jeholornis prima</i>	STM3-19	Avialae	Plant + Seed	Herbivory	6	0	2.77	97.23	0
<i>Jeholornis prima</i>	STM3-28	Avialae	Plant + Seed	Herbivory	18	0	1.74	98.26	0
<i>Jeholornis prima</i>	STM3-31	Avialae	Plant + Seed	Herbivory	43	0.02	0.94	98.99	0.05
<i>Limusaurus inextricabilis</i>	IVPP V15293	Theropoda	Plant + Seed	Herbivory/ Omnivory	185	0.44	3.94	63.62	32.01
<i>Limusaurus inextricabilis</i>	IVPP V15297	Theropoda	Plant + Seed	Herbivory/ Omnivory	201	0.15	21.87	45.56	32.43
<i>Psittacosaurus mazongshanensis</i>	IVPP V12165	Ornithischia	<i>Vertebrate</i>	Herbivory	55	0.53	5.78	2.02	91.67
<i>Psittacosaurus mongoliensis</i>	AMNH 6253	Ornithischia	<i>Vertebrate</i>	Herbivory	48	1.07	0.86	8.77	89.31
<i>Sinornithomimus dongi</i>	uncatalogued	Theropoda	Plant + Seed	Herbivory	1070	0.04	3.72	95.61	0.63
<i>Tarbosaurus bataar</i>	MPC-D 552-1	Theropoda	<i>Vertebrate</i>	Carnivory	682	5.76	2.1	8.77	83.36

The misidentification of Omnivores appears to be informative. For instance, the herbivory-dominated Omnivore (*Anas platyrhynchos*) was misidentified as a Plant and Seed feeder, while the vertivory-dominated Omnivore (*Calonectris leucomelas*) was mis-categorized as a Vertebrate Feeder. This implies that stomach stone shape does not just assign specimens to discrete categories, but rather captures the continuous nature of animal dietary and digestive ecology.

Even so, the models are not perfect, and several inherent factors likely contribute to mispredictions. One unavoidable factor is the unidirectional nature of stone abrasion; stomach stones only become more rounded over time, not rougher. Therefore, a stomach stone is expected to be rounded if it was rounded upon consumption, regardless of the diet or stomach muscularity

of the host. This likely explains instances where some Vertebrate Feeders and Invertebrate Feeders were misclassified as Plant and Seed Feeders (Supplementary Table S1). Additionally, our models do not account for seasonal dietary shifts, which are common in many birds (Billerman et al. 2022). The use of discrete dietary categories herein is a necessary simplification of this complex biological reality. For consistency and objectivity, we assigned specimens to these categories following the comprehensive framework of Wilman et al. (2014). This necessary simplification, combined with the lack of precise collection dates for some specimens, means that some of the observed intraspecific variation in stone shape (Supplementary Table S1) could reflect these dietary variations, potentially lowering the overall prediction accuracy.




Figure 3. Hypothetical evolutionary history of archosaur stomach muscularity, represented as a phylogenetic gradient map of the gastrolith shape proxy (principal component 1 [PC 1] value). Topology follows a composite phylogenetic framework assembled from published sources (see “Methods”; [Supplementary Data S4](#)) and is used as a scaffold for trait mapping. The circles next to the taxon names represent dentition status, and the absence of a circle represents a specimen without a known skull. Note that completely edentulous taxa generally have rounded gastroliths.

More broadly, the variation in gastrolith shape at the moment of ingestion—whether shaped by host animal selectivity, source environment lithology, or pre-ingestion transport history—is intrinsically unknown for both the extant and fossil specimens examined here, as rearing or capture locality ([Supplementary Table S6](#)) does not necessarily indicate where the gastroliths were acquired. However, if pre-consumption shape distribution were the primary driver of gastrolith roundness rather than intragastric abrasion, the diet–shape and muscularity–shape signals would not be expected to emerge with the clarity recovered here (78.57% diet accuracy; PC 1 vs. stomach muscularity $R^2 = 0.44$, $p < 0.001$) from such a heterogeneous extant sample. Pre-consumption variation therefore likely contributes to individual-level scatter—as invoked above for some misclassifications—but not to the broader diet–shape and muscularity–shape relationships reported here.

Despite these limitations and nuanced results, the models demonstrate predictive power far exceeding the 25% accuracy expected from random chance. The high accuracies achieved in our models strongly support the hypothesis that stomach stone shape contains a robust signal reflecting both archosaur diet and stomach muscularity. Therefore, while acknowledging its inherent uncertainties, this shape-based analysis serves as a potential new proxy for inferring archosaur dietary habits, to be used alongside classic methods such as osteology, stable isotopes, and gut content analysis. Such multiproxy approaches are increasingly advocated for robust paleodietary reconstruction in extinct taxa, and explicit frameworks developed for fossil birds provide a useful template for synthesizing independent lines of evidence ([Miller and Pittman 2021](#); [Miller et al. 2022](#)). The present gastrolith-shape analysis adds an independent proxy for gastrointestinal mechanical function (gastric abrasion capacity/stomach muscularity) that can be integrated alongside osteology, stable isotopes, and gut content evidence.

A practical road map for applying gastrolith shape information within a multiproxy framework begins with establishing the reliability of the stone assemblage as gastroliths. A conservative inclusion criterion is applied here, restricting scoring to stones preserved within, or immediately adjacent to, the abdominal region to

minimize inclusion of non-ingested clasts. Gastrolith shape is then evaluated using standardized criteria proposed by Best and Gionfriddo (1991). Analyses should preferentially focus on individuals with sufficient numbers of gastroliths (≥ 35 in the present dataset), although predictions based on smaller samples can also be reported, provided the associated uncertainty is explicitly acknowledged.

Once shape data are compiled, diet can be predicted using the trained classifier developed here, yielding coarse output categories expressed in the EltonTraits labels. These predictions are best treated as statistically-grounded hypotheses rather than definitive diet diagnoses, because the association between shape and diet is plausibly mediated by the mechanical environment of the stomach (details discussed later). Accordingly, interpretation should explicitly distinguish (1) the classifier’s dietary category prediction and (2) the mechanistic implication of the observed gastrolith morphotypes (e.g., whether stone rounding suggests sustained abrasion vs. limited abrasion), while recognizing that these two inferences are related but not identical. Finally, gastrolith-based predictions and mechanistic inferences should be evaluated alongside independent proxies (e.g., cranial and dental morphology, microwear, stable isotopes, gut contents). Concordance among independent lines of evidence strengthens dietary inference, whereas conflicts can be informative and should motivate targeted alternative hypotheses. In the following sections, gastrolith-shape-based predictions are compared with previous dietary interpretations derived from other proxies (primarily osteological evidence) to evaluate when this new proxy is most informative and what aspects of digestive function it most directly reflects.

Because fossil specimens preserving in situ stomach stones are rare, we report model outputs for all fossil individuals, including those with fewer than 35 stones, to maximize taxonomic coverage. However, predictions based on very small stone counts should be interpreted cautiously given our subsampling sensitivity analysis; consistent with this expectation, the two $n = 6$ specimens (*Iteravis*, IVPP V18958; and *Jeholornis*, STM 3-19) plot far outside the primary morphospace ([Supplementary Table S5](#)). In the specimens with at least 35 gastroliths, predicted diets are generally consistent

with previous interpretations in non-avian theropods (Table 1). By contrast, conflicts arise in Mesozoic birds, ornithischians, and sauropods, requiring reconciliation and motivating new hypotheses about their dietary habits and gastrointestinal properties. The Mesozoic birds *Bellulornis*, *Gansus*, and *Iteravis* have often been suggested to be herbivorous based on the presence of stomach stones (Y.-M. Wang et al. 2015; M. Wang et al. 2016), although more recently O'Connor (2019) inferred that *Gansus* and *Iteravis* are unlikely to be granivorous. O'Connor and Zhou (2019) further inferred omnivory in *Iteravis* based on the presence of fractured bones and carbonaceous remains reported in association with the specimen, although these remains are preserved external to the body cavity. The model developed in this study predicts these taxa as Vertebrate Feeders, which supports the more recent inference (O'Connor and Zhou 2019). The prediction is based on their angular stomach stones, which imply that their stomachs were not muscular enough to severely abrade them. Such a weakly muscular stomach would likely have been insufficient to process fibrous plant materials. *Iteravis* also has reduced dentition (Ju et al. 2020), which may indicate limited oral comminution. Therefore, the anatomical and stomach stone features of *Iteravis* and *Gansus* are consistent with a predominantly vertivorous lifestyle in these Mesozoic birds. In light of their high likelihood as Vertebrate Feeders (Table 1), the aquatic adaptations in the pelvis and hindlimbs of *Iteravis* and *Gansus* (You et al. 2006; Zhou et al. 2014; Wang et al. 2015) may further imply piscivory, a hypothesis that could be confirmed with future discovery of stomach contents. Unfortunately, the prediction of Vertebrate Feeding in *Bellulornis* cannot be checked against anatomical features until cranial remains are reported from this taxon.

The conflicts in the sauropods and ornithischians are less straightforward to reconcile, because both clades are very likely herbivorous based on osteology, yet gastrolith shape signals are not always concordant with those expectations. Sauropods have long been hypothesized to have performed limited oral processing and to have relied on prolonged retention and fermentation (Hummel and Clauss 2011). Consistent with this, a recently described *Diamantinasaurus* cololite provides direct evidence of herbivory and supports minimal oral processing, with digestion likely relying on fermentation and gut microflora (Poropat et al. 2025). Notably, no gastroliths were reported from the gut contents of *Diamantinasaurus*, consistent with the absence of a gastric mill suggested for sauropods (Wings and Sander 2007). Accordingly, where gastroliths are reported in sauropods, accidental ingestion during bulk feeding may remain a plausible explanation.

In contrast to sauropods, early-branching ornithischians generally have craniomandibular and dental specializations that indicate substantial oral processing. For example, the powerful adductor muscles and highly worn teeth of *Psittacosaurus* have been used to suggest their high-fiber diet (Serenó et al. 2010). Similarly, the basal ornithischians *Gasparinisaura* and *Haya* also have highly worn teeth and well-developed coronoid processes of the dentaries (Coria and Salgado 1996; Barta and Norell 2021), suggesting they also had a great deal of oral processing capability.

Taken together, these lines of evidence suggest that, at least in some herbivorous dinosaur clades, the stomach may not have been the primary site of mechanical breakdown. Accordingly, when gastroliths are present in basal ornithischians and sauropods, their shapes may not reflect sustained intragastric abrasion in the same way as in taxa with a strongly muscular stomach. Under this scenario, the stomachs of these two groups may have operated under a relatively low-abrasion regime (e.g., extant crocodylians,

carnivorous extant birds), although it is emphasized that this is a functional comparison rather than an anatomical one. Therefore, sauropod and ornithischian stomach stones would be likely to retain their original shapes at the time of ingestion. As such, conflicts between stomach stone-based predictions and previously inferred diets in ornithischians and sauropods likely reflect the evolutionary history of dinosaur stomachs and compensation by other organ systems. This reveals a nuance of our approach: while stomach stone shape is robust in assessing stomach muscularity among archosaurs, special care must be taken in using stomach stone shape to infer diet where other compensating food processing mechanisms are present. In this light, stomach stone shape may most accurately reflect diet in theropods, including birds, which lack extensive oral processing ability.

Nevertheless, the analyses presented here do not preclude alternative, non-exclusive functions of gastroliths in herbivorous ornithischians (e.g., geophagy and mineral supplementation or other physiological roles with potential metabolic significance). Standardized shape scoring for ornithischians remains limited in the present dataset (Table 1), which precludes robust inference about function based on the scored sample alone. Even so, the dominance of angular morphotypes in the scored ornithischians is not consistent with sustained intragastric abrasion expected under a muscular stomach. Dense abdominal gastrolith clusters are nevertheless reported repeatedly in *Psittacosaurus* (Osborn 1924; Sereno et al. 2010; Wang et al. 2026). Gastroliths have also been reported in additional ornithischian lineages, including *Haya* (Makovicky et al. 2011), *Changmiania* (Yang et al. 2020), *Gasparinisaura* (Cerdeña 2008), *Yinlong* (Xu et al. 2006), and *Tenontosaurus* (Nudds et al. 2022). The recent report of gastroliths in the early-branching pachycephalosaur *Zavacephale* (Chinzorig et al. 2025) further indicates that gastrolith occurrence was widespread among early-branching ornithischians. Recurrent occurrence across multiple individuals and lineages may suggest functional significance that does not depend on strong gastric abrasion. Distinguishing among alternative hypotheses will require broader taxonomic sampling and integration of independent proxies.

The ancestral-state reconstructions support a scenario in which muscular stomachs evolved in Theropoda, although future evaluations of gastroliths in a broader taxonomic sample might reveal independent origins of muscular stomachs. Indeed, some data show that stomach muscularity is evolutionarily plastic (Takasaki and Kobayashi 2024), and previous work shows that crocodilian and neornithine “gizzards” are not homologous (Takasaki and Kobayashi 2020b). It is also worth noting that gastroliths have also been reported in pterosaurs (Codorniu et al. 2013; Jiang et al. 2025), indicating that lithophagy occurred outside Dinosauria. However, comparable shape data are not yet available for reported pterosaur gastrolith assemblages, so their implications for ancestral ornithodiran stomach function remain uncertain. Nevertheless, the presently available data indicate that the muscular stomach originated at least within ancestral Maniraptoriformes, if it was not already present further back in the theropod lineage. As a phylogenetically independent comparison, *Limusaurus*, a highly unusual ceratosaur outside Maniraptoriformes, provides an informative functional case study (Xu et al. 2009; Wang et al. 2017). Through ontogeny, *Limusaurus* stopped replacing its teeth and became fully edentulous (Wang et al. 2017), while also acquiring moderately rounded stomach stones as an adult (Wang et al. 2017; Table 1). This suggests that, at least as adults, *Limusaurus* may have relied on a somewhat muscular, gastrolith-assisted stomach instead of teeth to break down plant material for digestion, but with less intensity than in

extant herbivorous birds (which have more rounded gastroliths). Given its unusual ontogenetic trajectory, this association may be unique to *Limusaurus* and therefore should not be used to infer the ancestral state of theropod stomach condition. Instead, *Limusaurus* may serve as a fossil example showing that reduced oral processing (edentulism) can co-occur with gastrolith-assisted gastric processing.

An early origin of a muscular stomach within Maniraptoriformes (and possibly deeper within Theropoda) potentially has broader implications for theropod evolution. Edentulism in theropods has long been considered an indicator of herbivory (Barrett 2005; Zanno and Makovicky 2011), but few studies have tried to explain what mechanisms substituted for oral processing. Our findings demonstrate that, where preserved, edentulous theropod taxa have highly rounded gastroliths indicative of enhanced gastrointestinal processing capabilities (Fig. 3). This suggests that the muscular stomach took up the role of the intense mechanical food processing system that is mandatory for herbivory in animals with high metabolic rates (Hummel et al. 2020). This is supported by the ontogenetic transition to edentulism and acquisition of moderately rounded stomach stones in *Limusaurus*, suggesting that, even within an individual lifetime, a muscularized stomach with gastroliths can effectively replace oral processing capabilities conferred by teeth. In this light, edentulism in theropods may not be an adaptation for herbivory itself, but a reflection of a transition to gastrointestinal food processing that released a selective constraint on the presence of teeth (Zheng et al. 2014; O'Connor 2019). Tooth loss may then have been advantageous in saving nutritional and energetic costs required for tooth replacement, which is estimated to have occurred every few months in dinosaurs (D'Emic et al. 2013, 2019). Thus, acquisition of a muscular stomach was likely an evolutionary innovation that enabled tooth reduction in many maniraptoriform groups, including ornithomimosaurs, therizinosaurs, oviraptorosaurs, and ultimately birds.

Offloading food processing to the muscular stomach may have had strong effects on theropod cranial constraints beyond enabling edentulism. In vertebrates, intensive oral processing is often associated with enlarged jaw adductor musculature, which occupies substantial cranial space and can constrain braincase organization and brain–endocranium disparity (Stedman et al. 2004; Penrose et al. 2016; Challands et al. 2020). Reduced reliance on oral processing in some theropods (e.g., ornithomimosaurs), potentially accompanied by reductions in these muscle groups (Cuff and Rayfield 2015), may therefore have relaxed the constraints and facilitated diversification of neurosensory systems along the avian stem lineage (Balanoff et al. 2013; Ksepka et al. 2020; Choiniere et al. 2021). Any implications for cognition remain speculative and are not directly tested here; cognitive performance in birds does not map simply onto absolute brain size, and neuronal packing densities and neuron numbers have been emphasized as additional correlates (e.g., Olkiewicz et al. 2016). Future comparative work integrating jaw muscle reconstructions, endocranial morphology, and sensory proxies across theropods will be necessary to evaluate whether shifts in gastrointestinal food processing contributed to patterns of avian cranial evolution.

Acknowledgments. The authors appreciate M. Eda (Hokkaido University Museum) and his volunteers for collecting and managing the specimens at the Hokkaido University Museum. F. Takaya (Botanic Garden, Hokkaido University) kindly allowed us to work on the specimens at the Botanic Garden, Hokkaido University. J. O'Connor kindly shared high-resolution photos of several Cretaceous bird gastroliths. We thank K. Chiba (Okayama University of Science), J. Yoshida (Fukushima Prefecture Museum), T. Tanaka (Hyogo Prefecture University), and M. Iijima (Institute of Vertebrate Paleontology) for

their helpful comments on earlier versions of the article. R.T. is funded by JSPS (17J06410, 20J01696, 23K13207). G.F.F. is funded by NSERC and the Vanier-Banting Commission.

Author Contribution. R.T. conceived and designed the study, collected and curated the data, performed all statistical, and morphometric analyses, wrote the analytical code, prepared the figures, acquired funding, and wrote the original draft. Y.K. supervised the project. A.R.F., T.C., and G.F.F. contributed to the review and editing of the manuscript. All authors read and approved the final version.

Competing Interests. The authors declare no competing interests

Data and Code Availability Statement. All data are provided as Supplementary Data S1–S4 and deposited with the Supplementary Material (Supplementary Figs. S1–S3 and Supplementary Tables S1–S6) at Dryad: <https://doi.org/10.5061/dryad.5mkkwh7k6>. The scripts used for the analyses are provided as Supplementary Data S5 and S6.

Literature Cited

- Balanoff, A. M., G. S. Bever, T. B. Rowe, and M. A. Norell. 2013. Evolutionary origins of the avian brain. *Nature* 501:93–96.
- Barlow, J., E. Klaas, and J. Lenz. 1963. Sunning of bank swallows and cliff swallows. *Condor* 65:438–440.
- Barrentine, C. D. 1980. The ingestion of grit by nestling barn swallows. *Journal of Field Ornithology* 51:368–371.
- Barrett, P. M. 2005. The diet of ostrich dinosaurs (Theropoda: Ornithomimosauria). *Palaeontology* 48:347–358.
- Barta, D. E., and M. A. Norell. 2021. The osteology of *Haya griva* (Dinosauria: Ornithischia) from the Late Cretaceous of Mongolia. *Bulletin of the American Museum of Natural History* 445:1–111.
- Beaune, D., C. Le Bohec, F. Lucas, M. Gauthier-Clerc, and Y. Le Maho. 2009. Stomach stones in king penguin chicks. *Polar Biology* 32:593–597.
- Best, L. B., and J. P. Gionfriddo. 1991. Characterization of grit use by cornfield birds. *The Wilson Bulletin* 103:68–82.
- Best, L. B., and J. P. Gionfriddo. 1994. Effects of surface texture and shape on grit selection by house sparrows and northern bobwhite. *The Wilson Bulletin* 106:689–695.
- Billerman, S. M., B. K. Keeney, P. G. Rodewald, and T. S. Schulenberg. 2022. *Birds of the world*. Cornell Laboratory of Ornithology, Ithaca, NY.
- Boyd, C. A. 2015. The systematic relationships and biogeographic history of ornithischian dinosaurs. *PeerJ* 3:e1523.
- Brown, C. R. 1976. Use of gravel by purple martins. *The Auk* 93:842–842.
- Cerda, I. A. 2008. Gastroliths in an ornithopod dinosaur. *Acta Palaeontologica Polonica* 53:351–355.
- Challands, T. J., J. D. Pardo, and A. M. Clement. 2020. Mandibular musculature constrains brain–endocranium disparity between sarcopterygians. *Royal Society Open Science* 7:200933.
- Chinzorig, T., R. Takasaki, J. Yoshida, R. T. Tucker, B. Buyantegsh, B. Mainbayar, K. Tsogtbaatar, and L. E. Zanno. 2025. A domed pachycephalosaur from the early Cretaceous of Mongolia. *Nature* 646:1138–1145.
- Choiniere, J. N., J. M. Neenan, L. Schmitz, D. P. Ford, K. E. J. Chapelle, A. M. Balanoff, J. S. Sipla, et al. 2021. Evolution of vision and hearing modalities in theropod dinosaurs. *Science* 372:610–613.
- Codorniu, L., L. M. Chiappe, and F. D. Cid. 2013. First occurrence of stomach stones in pterosaurs. *Journal of Vertebrate Paleontology* 33:647–654.
- Coria, R. A., and L. Salgado. 1996. A basal iguanodontian (Ornithischia: Ornithopoda) from the Late Cretaceous of South America. *Journal of Vertebrate Paleontology* 16:445–457.
- Cuff, A. R., and E. J. Rayfield. 2015. Retrodeformation and muscular reconstruction of ornithomimosaurian dinosaur crania. *PeerJ* 3:e1093.
- Dal Sasso, C., and S. Maganuco. 2011. *Scipionyx samniticus* (Theropoda: Compsognathidae) from the Lower Cretaceous of Italy. Osteology, ontogenetic assessment, phylogeny, soft tissue anatomy, taphonomy and paleobiology. *Memorie della Societa Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano* 37:1–281.

- D'Emic, M. D., J. A. Whitlock, K. M. Smith, D. C. Fisher, and J. A. Wilson. 2013. Evolution of high tooth replacement rates in sauropod dinosaurs. *PLoS ONE* 8:e69235.
- D'Emic, M. D., P. M. O'Connor, T. R. Pascucci, J. N. Gavras, E. Mardakhayava, and E. K. Lund. 2019. Evolution of high tooth replacement rates in theropod dinosaurs. *PLoS ONE* 14:e0224734.
- Fritz, J., J. Hummel, E. Kienle, O. Wings, W. J. Streich, and M. Clauss. 2011. Gizzard vs. teeth, it's a tie: food-processing efficiency in herbivorous birds and mammals and implications for dinosaur feeding strategies. *Paleobiology* 37:577–586.
- Gionfriddo, J. P., and L. B. Best. 1999. Grit use by birds: a review. Pp. 89–148 in V. Nolan Jr., E. D. Ketterson, and C. F. Thompson, eds. *Current ornithology*. Kluwer Academic/Plenum, New York.
- Hendrickx, C., S. A. Hartman, and O. Mateus. 2015. An overview of non-avian theropod discoveries and classification. *PalArch's Journal of Vertebrate Palaeontology* 12:1–73.
- Hummel, J., and M. Clauss. 2011. Sauropod feeding and digestive physiology. Pp. 11–32 in N. Klein, K. Remes, C. T. Gee, and P. M. Sander, eds. *Biology of the sauropod dinosaurs. Understanding the life of giants*. Indiana University Press, Bloomington.
- Hummel, J., M. Clauss, and K.-H. Südekum. 2020. Aspects of food comminution in ungulates and their consequences for energy budget. Pp. 87–101 in T. Martin, and W. v. Koenigswald, eds. *Mammalian teeth—form and function*. Friedrich Pfeil, Munich, Germany.
- Jenkinson, M. A., and R. M. Mengel. 1970. Ingestion of stones by goatsuckers (Caprimulgidae). *Condor* 72:236–237.
- Jiang, S., X. Zhang, Y. Wu, M. Zheng, A. W. A. Kellner, and X. Wang. 2025. First occurrence of phytoliths in pterosaurs-evidence for herbivory. *Science Bulletin* 70:3134–3138.
- Ju, S.-b., X.-r. Wang, Y.-c. Liu, and Y. Wang. 2020. A reassessment of *Iteravis huchzermeyeri* and *Gansus zheni* from the Jehol Biota in western Liaoning, China. *China Geology* 4:197–204.
- Kobayashi, Y., J.-C. Lu, Z.-M. Dong, R. Barsbold, Y. Azuma, and Y. Tomida. 1999. Herbivorous diet in an ornithomimid dinosaur. *Nature* 402:480–481.
- Ksepka, D. T., A. M. Balanoff, N. A. Smith, G. S. Bever, B. S. Bhullar, E. Bourdon, E. L. Braun, J. G. Burleigh, et al. 2020. Tempo and pattern of avian brain size evolution. *Current Biology* 30:2026–2036.
- Lambrecht, K. 1933. *Handbuch der palaeornithologie*. Gebrüder Borntraeger.
- Li, Z., Z. Zhou, M. Wang, and J. A. Clarke. 2014. A new specimen of large-bodied basal enantiornithine *Bohaiornis* from the early Cretaceous of China and the inference of feeding ecology in Mesozoic birds. *Journal of Paleontology* 88:99–108.
- Liu, S., Z. Li, A. M. Bailleul, M. Wang, and J. O'Connor. 2021. Investigating possible gastroliths in a referred specimen of *Bohaiornis guoi* (Aves: Enantiornithes). *Frontiers in Earth Science* 9:635727.
- Makovicky, P. J., B. M. Kilbourne, R. W. Sadleir, and M. A. Norell. 2011. A new basal ornithomimid (Dinosauria, Ornithischia) from the Late Cretaceous of Mongolia. *Journal of Vertebrate Paleontology* 31:626–640.
- McKeown, K. C. 1934. The food of birds from south-western New South Wales. *Records of the Australian Museum* 19:113–135.
- Miller, C. V., and M. Pittman. 2021. The diet of early birds based on modern and fossil evidence and a new framework for its reconstruction. *Biological Reviews* 96:2058–2112.
- Miller, C. V., M. Pittman, X. Wang, X. Zheng, and J. A. Bright. 2022. Diet of Mesozoic toothed birds (Longipterygidae) inferred from quantitative analysis of extant avian diet proxies. *BMC Biology* 20:101.
- Moore, S. J. 1998. The gizzard morphology of an avian herbivore: the domestic goose, *Anser anser*. *Australian Journal of Zoology* 46:345–357.
- Nudds, J. R., D. R. Lomax, and J. P. Tennant. 2022. Gastroliths and *Deinonychus* teeth associated with a skeleton of *Tenontosaurus* from the Cloverly Formation (Lower Cretaceous), Montana, USA. *Cretaceous Research* 140:105327.
- O'Connor, J. 2019. The trophic habits of early birds. *Palaeogeography, Palaeoclimatology, Palaeoecology* 513:178–195.
- O'Connor, J. K., and Z. Zhou. 2019. The evolution of the modern avian digestive system: insights from paravian fossils from the Yanliao and Jehol biotas. *Palaeontology* 63:13–27.
- Olkowicz, S., M. Kocourek, R. K. Lucan, M. Portes, W. T. Fitch, S. Herculano-Houzel, and P. Nemec. 2016. Birds have primate-like numbers of neurons in the forebrain. *Proceedings of the National Academy of Sciences USA* 113:7255–7260.
- Osborn, H. F. 1924. *Psittacosaurus and Protiguanodon: two Lower Cretaceous iguanodonts from Mongolia*. American Museum Novitates, no. 127. American Museum of Natural History, New York.
- Paradis, E., and K. Schliep. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* 35:526–528.
- Pennell, M. W., J. M. Eastman, G. J. Slater, J. W. Brown, J. C. Uyeda, R. G. FitzJohn, M. E. Alfaro, and L. J. Harmon. 2014. geiger v2.0: an expanded suite of methods for fitting macroevolutionary models to phylogenetic trees. *Bioinformatics* 30:2216–2218.
- Penrose, F., G. J. Kemp, and N. Jeffery. 2016. Scaling and accommodation of jaw adductor muscles in Canidae. *Anatomical Record* 299:951–966.
- Poropat, S. F., A. P. Tosolini, S. L. Beeston, M. J. Enchelmaier, A. H. Pentland, P. D. Mannion, P. Upchurch, et al. 2025. Fossilized gut contents elucidate the feeding habits of sauropod dinosaurs. *Current Biology* 35:2597–2613 e7.
- Potter, E. F. 1983. Ingestion of grit by wood warblers. *The Chat* 47:103–104.
- Powers, M. C. 1953. A new roundness scale for sedimentary particles. *Journal of Sedimentary Research* 23:117–119.
- R Core Team. 2020. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rörig, G. 1907. Magenuntersuchungen heimischer Raubvögel. *Arbeiten aus der Kaiserlichen Biologischen Anstalt für Land- und Forstwirtschaft* 5:237–265.
- Russell, R. D., and R. E. Taylor. 1937. Roundness and shape of Mississippi River sands. *Journal of Geology* 45:225–267.
- Sander, P. M., A. Christian, M. Clauss, R. Fechner, C. T. Gee, E. M. Griebeler, H. C. Gunga, et al. 2011. Biology of the sauropod dinosaurs: the evolution of gigantism. *Biological Reviews* 86:117–155.
- Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9:671–675.
- Sereno, P. C., Z. Xijin, and T. Lin. 2010. A new psittacosaur from Inner Mongolia and the parrot-like structure and function of the psittacosaur skull. *Proceedings of the Royal Society B* 277:199–209.
- Siegel-Causey, D. 1990. Gastroliths assist digestion in shags. *Notornis* 37:70–72.
- Stedman, H. H., B. W. Kozyak, A. Nelson, D. M. Thesier, L. T. Su, D. W. Low, C. R. Bridges, J. B. Shrager, N. Minugh-Purvis, and M. A. Mitchell. 2004. Myosin gene mutation correlates with anatomical changes in the human lineage. *Nature* 428:415–418.
- Takasaki, R., and Y. Kobayashi. 2020a. Effects of diet and gizzard muscularity on grit use in domestic chickens. *PeerJ* 8:e10277.
- Takasaki, R., and Y. Kobayashi. 2020b. Stomach histology of *Crocodylus siamensis* and *Gavialis gangeticus* reveals analogy of archosaur “gizzards,” with implication on crocodylian gastroliths function. *Acta Herpetologica* 15:111–118.
- Takasaki, R., and Y. Kobayashi. 2024. Anatomical description of neornithine stomach with implications on neornithine stomach morphology. *Journal of Anatomy* 245:787–796.
- Venables, W. N., and B. D. Ripley. 2013. *Modern applied statistics with S*. Springer Science & Business Media, New York.
- Wang, L., X. Li, Q. Wen, J. Chen, and R. Reisz. 2026. Gastroliths in hatchling *Psittacosaurus* show early dietary habits. *Science China Earth Sciences* 69:340–347.
- Wang, M., Z. Zhou, and S. Zhou. 2016. A new basal ornithuromorph bird (Aves: Ornithothoraces) from the Early Cretaceous of China with implication for morphology of early Ornithuromorpha. *Zoological Journal of the Linnean Society* 176:207–223.
- Wang, S., J. Stiegler, R. Amiot, X. Wang, G. H. Du, J. M. Clark, and X. Xu. 2017. Extreme ontogenetic changes in a ceratosaurian theropod. *Current Biology* 27:144–148.
- Wang, X., A. Cau, B. Guo, F. Ma, G. Qing, and Y. Liu. 2022. Intestinal preservation in a birdlike dinosaur supports conservatism in digestive canal evolution among theropods. *Scientific Reports* 12:19965.
- Wang, Y.-M., J. K. O'Connor, D.-Q. Li, and H.-L. You. 2015. New information on postcranial skeleton of the Early Cretaceous *Gansus yumenensis* (Aves: Ornithuromorpha). *Historical Biology* 28:666–679.

- Warton, D. I., R. A. Duursma, D. S. Falster, and S. Taskinen. 2012. smatr 3— an R package for estimation and inference about allometric lines. *Methods in Ecology and Evolution* 3:257–259.
- Wilman, H., J. Belmaker, J. Simpson, C. de la Rosa, M. M. Rivadeneira, and W. Jetz. 2014. EltonTraits 1.0: species-level foraging attributes of the world's birds and mammals. *Ecology* 95:2027.
- Wings, O. 2007. A review of gastrolith function with implications for fossil vertebrates and a revised classification. *Acta Palaeontologica Polonica* 52: 1–16.
- Wings, O., and P. M. Sander. 2007. No gastric mill in sauropod dinosaurs: new evidence from analysis of gastrolith mass and function in ostriches. *Proceedings of the Royal Society B* 274:635–640.
- Xu, X., C. A. Forster, J. M. Clark, and J. Mo. 2006. A basal ceratopsian with transitional features from the Late Jurassic of northwestern China. *Proceedings of the Royal Society B* 273:2135–2140.
- Xu, X., J. M. Clark, J. Mo, J. Choiniere, C. A. Forster, G. M. Erickson, D. W. Hone, et al. 2009. A Jurassic ceratopsian from China helps clarify avian digital homologies. *Nature* 459:940–944.
- Yang, Y., W. Wu, P.-E. Dieudonné, and P. Godefroit. 2020. A new basal ornithomimid dinosaur from the Lower Cretaceous of China. *PeerJ* 8:e9832.
- You, H. L., M. C. Lamanna, J. D. Harris, L. M. Chiappe, J. O'Connor, S. A. Ji, J. C. Lü, et al. 2006. A nearly modern amphibious bird from the Early Cretaceous of northwestern China. *Science* 312:1640–1643.
- Zanno, L. E., and P. J. Makovicky. 2011. Herbivorous ecomorphology and specialization patterns in theropod dinosaur evolution. *Proceedings of the National Academy of Sciences USA* 108:232–237.
- Zar, J. H. 2010. *Biostatistical analysis*. Prentice Hall, Englewood Cliffs, N.J.
- Zheng, X., J. K. O'Connor, F. Huchzermeyer, X. Wang, Y. Wang, X. Zhang, and Z. Zhou. 2014. New specimens of *Yanornis* indicate a piscivorous diet and modern alimentary canal. *PLoS ONE* 9:e95036.
- Zhou, S., J. K. O'Connor, and M. Wang. 2014. A new species from an ornithomimid (Aves: Ornithomimidae) dominated locality of the Jehol Biota. *Chinese Science Bulletin* 59:5366–5378.
- Zhou, Z., J. Clarke, F. Zhang, and O. Wings. 2004. Gastroliths in *Yanornis*: an indication of the earliest radical diet-switching and gizzard plasticity in the lineage leading to living birds? *Naturwissenschaften* 91:571–574.