

The origin of an invasive air sac system in sauropodomorph dinosaurs

Tito Aureliano^{1,2,3} | Aline M. Ghilardi² | Rodrigo T. Müller^{4,5} |
 Leonardo Kerber^{4,5} | Marcelo A. Fernandes³ | Fresia Ricardi-Branco¹ |
 Mathew J. Wedel⁶

¹Institute of Geosciences, University of Campinas (Unicamp), Campinas, Brazil

²Diversity, Ichnology and Osteohistology Laboratory (DINOlab), Department of Geology, Federal University of Rio Grande do Norte (UFRN), Natal, Brazil

³Laboratório de Paleocologia e Paleoecologia (LPP), Departamento de Ecologia e Biologia Evolutiva (DEBE), Federal University of São Carlos (UFSCar), São Carlos, Brazil

⁴Centro de Apoio à Pesquisa Paleontológica da Quarta Colônia (CAPPa), Federal University of Santa Maria (CAPPa/UFSM), São João do Polêsine, Brazil

⁵Programa de Pós-Graduação em Biodiversidade Animal, Federal University of Santa Maria (UFSM), Santa Maria, Brazil

⁶College of Osteopathic Medicine of the Pacific and College of Podiatric Medicine, Western University of Health Sciences, Pomona, USA

Correspondence

Tito Aureliano, Institute of Geosciences, University of Campinas (Unicamp), Campinas, California, Brazil.
 Email: aureliano.tito@gmail.com

Mathew J. Wedel, College of Osteopathic Medicine of the Pacific and College of Podiatric Medicine, Western University of Health Sciences, Pomona, USA.
 Email: mathew.wedel@gmail.com

Funding information

Fundaçao de Amparo à Pesquisa do Estado de São Paulo, Grant/Award Number: FAPESP 2019/16727-3;
 Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul, Grant/Award Numbers:
 21/2551-0000680-3, FAPERGS 1/2551-0002030-0; Conselho Nacional de Desenvolvimento Científico e Tecnológico, Grant/Award Numbers: 309414/2019-9, 422568/2018-0, 404095/2021-6, 307333/2021-3

Abstract

One of the most remarkable features in sauropod dinosaurs relates to their pneumatized skeletons permeated by a bird-like air sac system. Many studies described the late evolution and diversification of this trait in mid to late Mesozoic forms but few focused on the origin of the invasive respiratory diverticula in sauropodomorphs. Fortunately, it is possible to solve this thanks to the boom of new species described in the last decade as well as the broad accessibility of new technologies. Here we analyze the unaysaurid sauropodomorph *Macrocollum itaquii* from the Late Triassic (early Norian) of southern Brazil using micro-computed tomography. We describe the chronologically oldest and phylogenetically earliest unambiguous evidence of an invasive air sac system in a dinosaur. Surprisingly, this species presented a unique pattern of pneumatization in non-sauropod sauropodomorphs, with pneumatic foramina in posterior cervical and anterior dorsal vertebrae. This suggests that patterns of pneumatization were not cladistically consistent prior to the arrival of Jurassic eusauropods. Additionally, we describe the protocamerae tissue, a new type of pneumatic tissue with properties of both camellae and camerae. This reverts the previous hypothesis which stated that the skeletal pneumatization first evolved into camarae, and derived into delicate trabecular arrangements. This tissue is evidence of thin camellate-like tissue developing into larger chambers. Finally, *Macrocollum* is an example

of the gradual evolution of skeletal tissues responding to the fastly specializing Respiratory System of saurischian dinosaurs.

KEY WORDS

bone histology, ornithodira, respiratory system, South America, Triassic

1 | INTRODUCTION

Sauropods were an emblematic clade of long-necked dinosaurs that colonized all continents during the Jurassic and Cretaceous (Curry Rogers and Forster 2001; Cerda, Paulina Carabajal, et al., 2012; Ghilardi et al., 2016; Wiersma-Weyand et al., 2021; Rigby et al., 2021; Díez Díaz, 2022; Mo et al., 2023). Some lineages attained giant sizes, evolving into the largest animals to ever inhabit terrestrial landscapes (Carballido et al., 2017; González Riga et al., 2016; Lacovara et al., 2014; Otero et al., 2021). One of the features that allowed the body enlargement was a structural decrease of density caused by the presence of a bird-like air sac system (Perry et al., 2009; Sander et al., 2011). The identification of Postcranial Skeletal Pneumaticity (PSP) is the current method to identify the diverticular respiratory system in fossil taxa (Britt, 1994). The evaluation of unambiguous PSP requires the identification of deep vertebral fossae containing foramina connecting with internal pneumatic structures such as camerae and camellae (O'Connor, 2006). Unambiguous PSP has been extensively reported in neosauropods, especially titanosaurs (Wedel et al., 2000; Wedel, 2003; Schwarz & Fritsch, 2006; Woodward and Lehman 2009; Cerda, Salgado, & Powell, 2012; Zurriaguz & Powell, 2015; Bandeira et al., 2016). In contrast, few studies have focused on the origins of these diverticula within early sauropodomorphs, and much of the knowledge on non-sauropod sauropodomorphs is based on scarce evidence (Butler et al., 2012; Wedel, 2007, 2009; Yates et al., 2012). Recent evidence showing the absence of PSP in the earliest dinosaurs (Aureliano et al., 2022) suggested this trait was not homologous in ornithodirans and that invasive air sacs evolved at least three times independently, in theropods, sauropodomorphs, and pterosaurs. This discovery encouraged additional research into how and when these structures emerged during the Late Triassic. Here we evaluate the unaysaurid sauropodomorph *Macrocollum itaquii* (Müller, Langer, & Dias-da-Silva, 2018), the oldest and largest long-necked sauropodomorph of the lower Norian Candelária Sequence in South Brazil (Müller, Langer, & Dias-da-Silva, 2018). The micro-computed tomography of the skeleton of *Macrocollum* brings novel insights

into the gradual evolution of the Respiratory System in dinosaurs.

2 | MATERIALS AND METHODS

Institutional abbreviations: CAPPA/UFSM, Centro de Apoio à Pesquisa Paleontológica da Quarta Colônia, Universidade Federal de Santa Maria, São João do Polêsine, Rio Grande do Sul, Brazil; PVL, Paleovertebrate collection, Instituto “Miguel Lillo,” San Miguel de Tucumán, Salta, Argentina; NHM, Natural History Museum, London, United Kingdom; ULBRA, Centro de Apoio à Pesquisa Paleontológica da Quarta Colônia, Universidade Federal de Santa Maria, São João do Polêsine, Rio Grande do Sul, Brazil (previously Museu de Ciências Naturais, Universidade Luterana do Brasil, Canoas, Brazil).

2.1 | Specimen

The studied specimen corresponds to a paratype of *Macrocollum itaquii* (CAPPA/UFSM 0001b), an almost complete articulated skeleton of an unaysaurid sauropodomorph (Müller, Langer, & Dias-da-Silva, 2018). This specimen was found associated with two other skeletons (holotype and paratype) resulting in the oldest evidence of gregarious behavior in dinosaurs. There are yet no histological thin sections to address the skeletochronology of this specimen, but the scapula and the coracoid are partially unfused. Some vertebrae present fused neurocentral sutures to a varied degree along the column, some neural arches remain not entirely merged, which suggests a subadult ontogenetic stage (Brochu, 1996; Müller et al., 2015).

2.2 | Locality and horizon

Macrocollum specimens were excavated at the Wachholz Site ($29^{\circ}36'46.42''S$; $53^{\circ}15'54.06''W$), Agudo municipality, Rio Grande do Sul state, South Brazil (Müller et al., 2015). This site is part of the Upper Triassic (lower Norian) Candelária Sequence, Paraná Basin. Chronocorrelated strata were dated at 225.42 ± 0.37 Ma (Langer et al., 2018).




FIGURE 1 Skeletal reconstruction of the unaysaurid sauropodomorph *Macrocollum* (CAPPA/UFSM 0001b) showing vertebral elements along the spine and putative reconstruction of the air sac systems involved. (a) Pneumatic posterior cervical vertebra and a cross-section CT slice in b. (c) a pneumatized anterior dorsal vertebra with cross-section CT slice in d, and detail of the pneumatic foramen in e. (f) Detail of the pneumatic foramen in a reconstructed 3D model of the element. (g) Anterior cervical element (apneumatic). (h) Posterior dorsal vertebra shows no traces of PSP. The sacral series (i), as well as the anterior (k) and mid-caudal (j) series are apneumatic. a, g, h, j, and k are in left lateral view. c, e and f are in right lateral view. i is in dorsal view. ABD, abdominal diverticula; CER, cervical diverticula; LUN, lung; pf, pneumatic foramen. The reconstruction was made by Rodrigo T. Müller. Scale bar of the skeletal reconstruction = 500 mm; a-j = 20 mm.

2.3 | Taphonomic remarks

The CAPPA/UFSM 0001 series of specimens were moderately affected by taphonomic compression, showing both subtle diagenetic distortions and cracks (Müller, Garcia, et al., 2018). However, the external overall morphology is well preserved, except for smaller foramina, which will be further explained in the results section.

2.4 | Anatomical nomenclature and terminology

We apply Wilson's terms to describe vertebral laminae, fossae, and their landmarks (Wilson, 1999, 2012; Wilson

et al., 2011), Wedel's for vertebral pneumaticity (Wedel, 2003, 2007; Wedel et al., 2000), and O'Connor's method to evaluate unambiguous PSP (O'Connor, 2006).

2.5 | Microtomography (μ CT scan)

A posterior (eighth) cervical vertebra and an anterior (second) dorsal vertebra of *Macrocollum* were scanned in a Bruker-Skyscan 1,173 microtomographer (130 kV μ -focus X-ray source; voxel size = 0.15 mm). This μ CT scanner is located at the *Instituto do Petróleo e dos Recursos Naturais*, Universidade Católica do Rio Grande do Sul/PUCRS), Porto Alegre, Brazil. We used the open-source software 3D-Slicer v5.2 (Fedorov et al., 2012) and




FIGURE 2 Micro-CT scan of the posterior (eighth) cervical vertebra of the unaysaurid sauropodomorph *Macrocollum* (CAPPA/UFSCM 0001b) in cross-section (a–k) and lateral views (l, m). (a–k) presents a cross-section series from posterior to anterior-most positions, respectively. (l) and (m) show a narrow longitudinally-elongated chamber in the centrum and a decrease in trabecular density in the prezygapophysis and the neural spine. c, centrum; ccv, a chamber in the centrum; cdf, centrodiapophyseal fossa; cdl, centrodiapophyseal lamina; cpof, centropostzygapophyseal fossa; d, diapophysis; nc, neural canal; ncf, neural canal foramen; pf, pneumatic foramen; po, postzygapophysis; pocdf, postzygapophyseal centrodiapophyseal fossa; pr, prezygapophysis; prcdf, prezygapophyseal centrodiapophyseal fossa; ptc, protocamera; s, neural spine/ spof, spinopostzygapophyseal fossa. Scale bar = 10 mm.

CloudCompare v2.12.4 (CloudCompare, 2018) to process the data, to generate 3D reconstructions, and to apply density-based color grades based on bone tissues (Aureliano et al., 2020). *ImageJ* v1.52 (Schneider et al., 2012) was used for digital measurements. All microtomography data were uploaded to the Morphobank platform and is available through this link: <http://morphobank.org/permalink/?P4526>.

3 | RESULTS

We analyzed the vertebrae of the sauropodomorph *Macrocollum itaquii* employing both macroscopic inspection and computed tomography (Figure 1). The vertebral




FIGURE 3 Micro-CT scan of the anterior (second) dorsal vertebra of the unaysaurid sauropodomorph *Macrocollum* (CAPPA/UFSCM 0001b). (a) and (b) show cross-sections of the entire vertebra in anterior view at the approximate midpoint. (e) and (f) show midshaft slices in lateral view. (f) shows three fossae in the neural arch (cprf, cdf and cpof). c, centrum; cdf, centrodiapophyseal fossa; cdl, centrodiapophyseal lamina; ctr, chaotic trabeculae; cpof, centropostzygapophyseal fossa; cpol, centropostzygapophyseal lamina; cprf, centroprezygapophyseal fossa; d, diapophysis; dia, diagenetic artifact; nc, neural canal; ncf, neural canal foramen; pf, pneumatic foramen; po, postzygapophysis; pocdf, postzygapophyseal centrodiapophyseal fossa; pr, prezygapophysis; prcdf, prezygapophyseal centrodiapophyseal fossa; ptc, protocamera; s, neural spine. Scale bar = 10 mm.

tissue showed moderate evidence of diagenetic effects, mainly in the form of cracking and mineral infills, which altered the bone morphology slightly. This condition made it difficult to detect smaller nutritional foramina but, fortunately, did not affect the preservation of the larger pneumatic ones. It was thus possible to detect evidence of unambiguous PSP, which varies along the axial skeleton. Only presacral vertebrae demonstrated the presence of pneumatic foramina. The µCT scan of the posterior cervical element (Figure 2) revealed that the centrum comprises mostly dense trabecular bone except for a narrow longitudinally-elongated chamber connecting with two lateral pneumatic foramina. There is also a decrease in trabecular density in the neural arch. Several fossae (spof, cpof and cdf) connect with internal pneumatic

TABLE 1 Evaluation of postcranial skeletal pneumaticity in sauropodomorphs and the herrerasaurid *Gnathovorax*. Question marks indicate unknown information and/or elements.

Taxon/specimen	Age	Anterior cervical	Posterior cervical	Anterior dorsal	Posterior dorsal	Sacral	Anterior caudal	Reference
<i>Gnathovorax</i> (CAPPA/UFSM 0009)	Carnian	No	No	No	No	No	No	(Aureliano et al., 2022)
<i>Buriolestes</i> (CAPPA/UFSM 0035)	Carnian	No	No	No	No	No	No	(Aureliano et al., 2022)
<i>Pampadromaeus</i> (ULBRAPV016)	Carnian	?	?	No	?	?	?	(Aureliano et al., 2022)
<i>Macrocollum</i> (CAPPA/UFSM 0001b)	early Norian	No	Yes	Yes	No	No	No	This work
<i>Plateosaurus</i> (AMNH 6810)	late Norian - Rhaetian	No	Yes	No	No	No	No	Yates et al. (2012)
<i>Eucnemesaurus</i> (BP/1/6107)	late Norian - Rhaetian	?	?	?	Yes	?	?	Yates et al. (2012)
<i>Aardonyx</i> (BP/1/6566)	Hettangian	No	No	No	Yes	Yes	No	Yates et al. (2012)
<i>Antenoritrus</i> (BP/1/4952)	Sinemurian - Pliensbachian	?	?	No	Yes	?	No	Yates et al. (2012)
<i>Haplocaelosaurus</i> (CM879)	Kimmeridgian	Yes	Yes	Yes	Yes	Yes	Yes	Wedel (2009)
<i>Saltasaurus</i> (PVL 4017–214, 47, 192)	Maastrichtian	Yes	Yes	Yes	Yes	Yes	Yes	(Cerda, Salgado, & Powell, 2012); Zurriaguz and Powell (2015)




FIGURE 4 Chronological and phylogenetic evolution of postcranial skeletal pneumaticity in sauropodomorph dinosaurs, and the herrerasaurid *Gnathovorax* as an outgroup. 1, Sauropodomorpha. 2, Sauropoda. Based on the topology by Yates et al. (2012) and Müller, Langer, and Dias-da-Silva (2018). Silhouettes are from [Phylopic.org](#) by Bruno Navarro, Mathew Wedel, Ryan Soledade, and Scott Hartman.

chambers. These chambers are neither camerata nor camellata, but a new type of tissue with an intermediate texture. We propose the term “protocamerata chambers” (ptc) for these structures which are not large enough to be considered camerae, but also present a camellata array internally. The µCT scan of the anterior dorsal vertebra (Figure 3), on the other hand, shows only chaotic apneumatic trabeculae in the centrum with no lateral pneumatic foramina. Only the neural arch is pneumatized, encompassing well-defined laminae connecting with the internal protocamerata tissue (ptc) throughout the fossae (cdf, pocdf, and prcdf). The protocamerata tissue also expands ventrodorsally to the interior base of the spinal arch.

4 | DISCUSSION

An evaluation of the herrerasaurid *Gnathovorax* and the early sauropodomorphs *Buriolestes* and *Pampadromaeus* found evidence suggesting the absence of PSP in early saurischians during the Carnian (Aureliano et al., 2022). We analyzed the unaysaurid *Macrocollum* from the early Norian of South Brazil. The presence of well-developed fossae connected with internal chambers throughout foramina is the definitive evidence of unambiguous PSP (O'Connor, 2006). Consequently, this find represents the oldest evidence of an invasive air sac system in Sauropodomorpha.

Previously, Yates et al. (2012) analyzed the patterns of pneumatization in Rhaetian and Early Jurassic sauropodomorphs, finding PSP to be restricted to the posterior dorsal elements, as in *Eucnemesaurus*, *Aardonyx*, and *Antenonitrus* (Yates et al., 2012). This evidence led the authors to suggest that the pneumatization process in early forms was influenced directly and solely by the abdominal air sac system, a term defined in previous studies on extant birds (McLelland, 1989). According to their hypothesis, only in eusauropods would the cervical air sac system develop with phyletic consistency, and not sporadically as in non-gravisaurian taxa. The presence of pneumatic foramina invading the centrum in a posterior cervical vertebra and not in an anterior dorsal element in *Macrocollum* is intriguing and so far unique in early sauropodomorphs (see Table 1). The neural arches of *Macrocollum* also showed many connections between the interior pneumatic architecture and the outer fossae (cdf, cprf, cpof; see Figures 2 and 3). Some *Plateosaurus* specimens were also interpreted as showing limited and restricted cervical pneumatization (e.g., AMNH 6810) and were thus considered exceptions in the evolution of the group (Yates et al., 2012). *Pantydraco* could potentially be another exception (Yates et al., 2012) but the absence of almost the entire sacral and dorsal series precludes unequivocal determination of PSP features in this taxon for the time being.

The presence of both cervical and anterior dorsal pneumatization in *Macrocollum* is unique within non-gravisaurians, and could suggest the influence of both cervical and lung air sac systems invading the vertebral series. However, the absence of pneumatic foramina in the dorsal centra might be indicative that lung and abdominal diverticula were not invading the skeleton. The absence of evidence of PSP in posterior dorsal elements in chronologically and phylogenetically earlier forms prior to *Eucnemesaurus* contrasts with the pattern of pneumatization observed in later forms, specially anchisaurians (see Table 1 and Figure 4). The evidence of unambiguous PSP in *Macrocollum*, *Pantydraco*, and *Plateosaurus*, when compared with the apneumatic skeleton of *Buriolestes*, *Pampadromaeus*, *Thecodontosaurus*, *Efraasia*, and *Massospondylus*, demonstrates that the initial evolution of diverticular ventilation systems did not follow a pronounced pattern (at least at our current phylogenetic resolution).

Macrocollum also presented a new type of pneumatic tissue, showing properties of both camerae and camellae as defined by Wedel et al. (2000). We named this tissue “protocamerae” (see ptc in Figures 2 and 3). This evidence supports a previous hypothesis from Aureliano et al. (2022) in which the pneumatic tissues would not have evolved as camerae in the Jurassic and then camellae in the Cretaceous but actually derived from delicate camellata-like trabeculae. The structural change from the apneumatic pseudo-polycamerata tissue in *Pampadromaeus* during the Carnian (Aureliano et al., 2022) to the protocamerae in *Macrocollum* demonstrates the possible developmental trajectory of PSP during the earliest evolution of sauropodomorph dinosaurs. Additionally, Yates et al. (2012) observed that the pneumatization throughout the infradiapophyseal fossae appears later in neosauropods due to the presence of camellata tissue, which would increase the volume and the rigidity of the neural arch, allowing the invasion of the diverticula. The invasion of air sacs throughout the cdf, cprf and cpof in *Macrocollum* could have been facilitated by the presence of this protocamerata tissue, with a similar structural consequence.

5 | CONCLUSIONS

The evidence of unambiguous postcranial pneumaticity in the unaysaurid *Macrocollum itaquii* from the early Norian of Southern Brazil resulted in several insights into the evolution of the air sac systems. Some of the highlights are listed below:

- The chronologically oldest and phylogenetically earliest evidence described to date of an invasive air sac system in a dinosaur.

- The pattern of pneumatization seen in the posterior cervical and anterior dorsal vertebrae of *Macrocollum* is unique in non-sauropod sauropodomorphs. Our data support previous hypotheses that this clade did not develop a consistent and extensive pattern of PSP until the evolution of eusauropods in the Jurassic.
- We described a new type of pneumatic tissue, the protocamerae, showing properties of both camellate and camerata tissues found later in the Mesozoic. This changes the previous hypothesis which stated that the pneumatization of the skeleton first evolved into camarae, and only later derived into delicate trabecular camellae. The protocamerata tissue is evidence of delicate thin tissue forming larger chambers.

Macrocollum is an example of gradual evolution of both morphology and histology, with the skeletal tissues adapting to the invasion of the air sac diverticula. We suggest that the great number of new taxa and specimens of Late Triassic sauropodomorphs described in the recent decade should be also sampled since they have the potential to provide a high-resolution understanding of the evolution of the Respiratory System in dinosaurs. Finally, the usage of microtomography coupled with histological thin sections should be encouraged in future approaches to achieve this goal.

AUTHOR CONTRIBUTIONS

Tito Aureliano: Conceptualization; investigation; writing – original draft; methodology; validation; visualization; writing – review and editing; software; formal analysis; project administration; data curation. **Aline M. Ghilardi:** Conceptualization; investigation; methodology; validation; software; formal analysis; data curation; supervision. **Rodrigo T. Müller:** Writing – original draft; writing – review and editing; data curation; supervision; resources; validation. **Leonardo Kerber:** Investigation; validation; writing – review and editing; project administration; supervision; data curation; resources. **Marcelo A. Fernandes:** Supervision; data curation; resources; writing – review and editing; validation; project administration. **Fresia Ricardi-Branco:** Funding acquisition; writing – original draft; writing – review and editing; validation; formal analysis; project administration; supervision; data curation; resources. **Mathew J. Wedel:** Conceptualization; investigation; funding acquisition; writing – original draft; writing – review and editing; validation; methodology; formal analysis; supervision; data curation.

ACKNOWLEDGMENTS

We thank Adolpho Herbert Augustin for conducting the microtomography. We also would like to thank Pedro

Buck, Felipe Pinheiro, Ariel Martine, Giorgio Basilici, Blair McPhee, and Ignacio Cerdá for their comments on an earlier version of this article. Research grants, scholarships, and financial aid were provided by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP 2019/16727-3) “Taphonomical landscapes” (Fresia Ricardi-Branco), Fundação de Amparo à Pesquisa do Estado do Rio Grande do Sul (FAPERGS 1/2551-0002030-0, Leonardo Kerber; 21/2551-0000680-3, Rodrigo T. Müller), and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq grant 307333/2021-3, Fresia Ricardi-Branco; CNPq 404095/2021-6, Rodrigo T. Müller; 422568/2018-0; 309414/2019-9, Leonardo Kerber).

CONFLICT OF INTEREST STATEMENT

The authors declare no competing interests.

DATA AVAILABILITY STATEMENT

All fossils are housed in a public research institution and can be accessed upon request and addressed to the collection's curator. Microtomography data and specimen photographs are available in a Morphobank project linked to this article.

ORCID

Tito Aureliano  <https://orcid.org/0000-0002-6918-0519>
Rodrigo T. Müller  <https://orcid.org/0000-0001-8894-9875>

REFERENCES

- Aureliano, T., Ghilardi, A. M., Müller, R. T., Kerber, L., Pretto, F. A., Fernandes, M. A., Ricardi-Branco, F., & Wedel, M. J. (2022). The absence of an invasive air sac system in the earliest dinosaurs suggests multiple origins of vertebral pneumaticity. *Scientific Reports*, 12, 20844.
- Aureliano, T., Ghilardi, A. M., Silva-Junior, J. C. G., Martinelli, A. G., Ribeiro, L. C. B., Marinho, T., Fernandes, M. A., Ricardi-Branco, F., & Sander, P. M. (2020). Influence of taphonomy on histological evidence for vertebral pneumaticity in an upper cretaceous titanosaur from South America. *Cretaceous Research*, 108, 104337.
- Bandeira, K. L. N., Simbras, F. M., Machado, E. B., et al. (2016). A new giant Titanosauria (Dinosauria: Sauropoda) from the late cretaceous Bauru group. *Brazil. PLoS One*, 11, e0163373.
- Britt, B. B. (1994). Pneumatic postcranial bones in dinosaurs and other archosaurs.
- Brochu, C. A. (1996). Closure of neurocentral sutures during crocodilian ontogeny: Implications for maturity assessment in fossil archosaurs. *Journal of Vertebrate Paleontology*, 16, 49–62.
- Butler, R. J., Barrett, P. M., & Gower, D. J. (2012). Reassessment of the evidence for postcranial skeletal pneumaticity in Triassic archosaurs, and the early evolution of the avian respiratory system. *PLoS One*, 7, e34094.
- Carballido, J. L., Pol, D., Otero, A., Cerdá, I. A., Salgado, L., Garrido, A. C., Ramezani, J., Cúneo, N. R., & Krause, J. M.

- (2017). A new giant titanosaur sheds light on body mass evolution among sauropod dinosaurs. *Proceedings of the Royal Society B*, 284, 20171219.
- Cerda, I. A., Paulina Carabajal, A., Salgado, L., Coria, R. A., Reguero, M. A., Tambussi, C. P., & Moly, J. J. (2012). The first record of a sauropod dinosaur from Antarctica. *Naturwissenschaften*, 99, 83–87.
- Cerda, I. A., Salgado, L., & Powell, J. E. (2012). Extreme postcranial pneumaticity in sauropod dinosaurs from South America. *Paläontologische Zeitschrift*, 86, 441–449.
- CloudCompare. (2018). Version 2.9. 1 [GPL software].
- Curry Rogers, K., & Forster, C. A. (2001). The last of the dinosaur titans: A new sauropod from Madagascar. *Nature*, 412, 530–534.
- Díez Díaz, V. (2022). Titanosaur boom. *Nature Ecology & Evolution*, 6, 251–252.
- Fedorov, A., Beichel, R., Kalpathy-Cramer, J., Finet, J., Fillion-Robin, J. C., Pujol, S., Bauer, C., Jennings, D., Fennessy, F., Sonka, M., Buatti, J., Aylward, S., Miller, J. V., Pieper, S., & Kikinis, R. (2012). 3D slicer as an image computing platform for the quantitative imaging network. *Magnetic Resonance Imaging*, 30, 1323–1341.
- Ghilardi, A. M., Aureliano, T., Duque, R. R. C., Fernandes, M. A., Barreto, A. M. F., & Chinsamy, A. (2016). A new titanosaur from the lower cretaceous of Brazil. *Cretaceous Research*, 67, 16–24.
- González Riga, B. J., Lamanna, M. C., Ortiz David, L. D., Calvo, J. O., & Coria, J. P. (2016). A gigantic new dinosaur from Argentina and the evolution of the sauropod hind foot. *Scientific Reports*, 6, 19165.
- Lacovara, K. J., Lamanna, M. C., Ibáñez, L. M., Poole, J. C., Schroeter, E. R., Ullmann, P. V., Voegeli, K. K., Boles, Z. M., Carter, A. M., Fowler, E. K., Egerton, V. M., Moyer, A. E., Coughenour, C. L., Schein, J. P., Harris, J. D., Martínez, R. D., & Novas, F. E. (2014). A gigantic, exceptionally complete titanosaurian sauropod dinosaur from southern Patagonia Argentina. *Scientific Reports*, 4, 6196.
- Langer, M. C., Ramezani, J., & Da Rosa, Á. A. S. (2018). U-Pb age constraints on dinosaur rise from South Brazil. *Gondwana Research*, 57, 133–140.
- McLellan, J. (1989). *Anatomy of the lungs and air sacs. Form and function in birds*, 4, 221–279.
- Mo, J., Ma, F., Yu, Y., & Xu, X. (2023). A new Titanosauriform sauropod with an unusual tail from the lower cretaceous of northeastern China. *Cretaceous Research*, 144, 105449.
- Müller, R. T., da Rosa, Á. A. S., Roberto da Silva, L., Aires, A. S. S., Pacheco, C. P., Pavanatto, A. E. B., & Dias-da-Silva, S. (2015). Wachholz, a new exquisite dinosaur-bearing fossiliferous site from the upper Triassic of southern Brazil. *Journal of South American Earth Sciences*, 61, 120–128.
- Müller, R. T., Garcia, M. S., Da-Rosa, Á. A. S., & Dias-da-Silva, S. (2018). Under pressure: Effect of sedimentary compression on the iliac morphology of early sauropodomorphs. *Journal of South American Earth Sciences*, 88, 345–351.
- Müller, R. T., Langer, M. C., & Dias-da-Silva, S. (2018). An exceptionally preserved association of complete dinosaur skeletons reveals the oldest long-necked sauropodomorphs. *Biology Letters*, 14, 20180633. <https://doi.org/10.1098/rsbl.2018.0633>
- O'Connor, P. M. (2006). Postcranial pneumaticity: An evaluation of soft-tissue influences on the postcranial skeleton and the reconstruction of pulmonary anatomy in archosaurs. *Journal of Morphology*, 267, 1199–1226.
- Otero, A., Carballido, J. L., Salgado, L., Canudo, J. I., & Garrido, A. C. (2021). Report of a giant titanosaur sauropod from the upper cretaceous of Neuquén Province Argentina. *Cretaceous Research*, 122, 104754.
- Perry, S. F., Christian, A., Breuer, T., et al. (2009). Implications of an avian-style respiratory system for gigantism in sauropod dinosaurs. *Journal of Experimental Zoology. Part A, Ecological Genetics and Physiology*, 311, 600–610.
- Rigby, S. L., Poropat, S. F., Mannion, P. D., Pentland, A. H., Sloan, T., Rumbold, S. J., Webster, C. B., & Elliott, D. A. (2021). A juvenile *Diamantinasaurus matildae* (Dinosauria: Titanosauria) from the upper cretaceous Winton formation of Queensland, Australia, with implications for sauropod ontogeny. *Journal of Vertebrate Paleontology*, 41(6), p.e2047991. <https://doi.org/10.1080/02724634.2021.2047991>
- Sander, P. M., Christian, A., Clauss, M., Fechner, R., Gee, C. T., Griebeler, E. M., Gunga, H. C., Hummel, J., Mallison, H., Perry, S. F., Preuschoft, H., Rauhut, O. W. M., Remes, K., Tütken, T., Wings, O., & Witzel, U. (2011). Biology of the sauropod dinosaurs: The evolution of gigantism. *Biological Reviews of the Cambridge Philosophical Society*, 86, 117–155.
- Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH image to ImageJ: 25 years of image analysis. *Nature Methods*, 9, 671–675.
- Schwarz, D., & Fritsch, G. (2006). Pneumatic structures in the cervical vertebrae of the late Jurassic Tendaguru sauropods *brachiosaurus brancai* and *Dicraeosaurus*. *Elogiae Geologicae Helveticae*, 99, 65–78.
- Wedel, M. J. (2003). Vertebral pneumaticity, air sacs, and the physiology of sauropod dinosaurs. *Paleobiology*, 29, 243–255.
- Wedel, M. J. (2007). What pneumaticity tells us about “prosauropods”, and vice versa. *Special Papers in Palaeontology*, 77, 207–222.
- Wedel, M. J. (2009). Evidence for bird-like air sacs in saurischian dinosaurs. *Journal of Experimental Zoology. Part A, Ecological Genetics and Physiology*, 311, 611–628.
- Wedel, M. J., Cifelli, R. L., & Sanders, R. K. (2000). Osteology, paleobiology, and relationships of the sauropod dinosaur *Sauroposeidon*. *Acta Palaeontologica Polonica*, 45(4), 343.
- Wiersma-Weyand, K., Canoville, A., Siber, H.-J., & Sander, P. M. (2021). Testing hypothesis of skeletal unity using bone histology: The case of the sauropod remains from the Howe-Stephens and Howe Scott quarries. *Palaeontol Electronica*. <https://doi.org/10.26879/766>
- Wilson, J. A. (1999). A nomenclature for vertebral laminae in sauropods and other saurischian dinosaurs. *Journal of Vertebrate Paleontology*, 19, 639–653.
- Wilson, J. A. (2012). New vertebral laminae and patterns of serial variation in vertebral laminae of sauropod dinosaurs.
- Wilson, J. A., D'Emic, M. D., Ikejiri, T., et al. (2011). A nomenclature for vertebral fossae in sauropods and other saurischian dinosaurs. *PLoS One*, 6, e17114.
- Woodward, H. N., & Lehman, T. M. (2009). Bone histology and microanatomy of *Alamosaurus sanjuanensis* (Sauropoda:

- Titanosauria) from the Maastrichtian of big bend National Park, Texas. *Journal of Vertebrate Paleontology*, 29, 807–821.
- Yates, A. M., Wedel, M. J., & Bonnan, M. F. (2012). The early evolution of postcranial skeletal Pneumaticity in sauropodomorph dinosaurs. *Acta Palaeontologica Polonica*, 57, 85–100.
- Zurriaguz, V., & Powell, J. E. (2015). New contributions to the pre-sacral osteology of *Saltasaurus loricatus* (Sauropoda, Titanosauria) from the upper cretaceous of northern Argentina. *Cretaceous Research*, 54, 283–300.

How to cite this article: Aureliano, T., Ghilardi, A. M., Müller, R. T., Kerber, L., Fernandes, M. A., Ricardi-Branco, F., & Wedel, M. J. (2023). The origin of an invasive air sac system in sauropodomorph dinosaurs. *The Anatomical Record*, 1–9. <https://doi.org/10.1002/ar.25209>